

*tinillus validus* (Horn) in North America. *Annals Amer. Ent. Soc.* 48:417-421.

RILEY, C. V.

1892. Systematic relations of *Platypsyllus*, as determined by the larva. In *Castorologia*, H. T. Martin. Wm. Drysdale and Co., Montreal. xvi plus 238 pp.

RITSEMA, J.

1869. Petites nouvelles. *Petites nouvelles entomologiques*. Sept. 15, 1rst. year.

NOTES AND SYNONYMY OF SOME NEOTROPICAL  
SPHEX AND ISODONTIA DESCRIBED BY  
E. TASCHENBERG AND S. ROHWER

(Hymenoptera: Sphecidae)

A. S. MENKE

*University of California, Davis*

The following notes are based primarily on Taschenberg type material generously lent by J. O. Husing of the Zoologisches Institut, Martin-Luther-Universität, Halle, East Germany. In addition, a recent visit to the United States National Museum allowed the author to examine the Rohwer types in that Institution. I am designating lectotypes for the following Taschenberg species: *Isodontia nigrocoerulea*, *Sphex argentinus* and *micans*.

ISODONTIA CYANIPENNIS (Fabricius)

*Sphex cyanipennis* Fabricius, 1793. *Ent. Syst.*, 2:200 (Cayenne, French Guiana, type lost).

*Sphex nigrocoerulea* Taschenberg, 1869. *Zeitsch. Ges. Naturw. Halle*, 34:415 (Lectotype ♀ and paralectotype ♀, Venezuela, present designation).

*Isodontia bipunctata* Rohwer, 1913. *Proc. U.S. Nat. Mus.*, 44:452 (holotype ♀, Canal Zone, Panama). *New synonymy*.

J. van der Vecht (1961) suggested that Fabricius' *cyanipennis* is the same species described by Taschenberg as *nigrocoerulea*. However, Richards (1937) described a metallic blue *Isodontia* from a male collected in British Guiana, which may be a synonym of *cyanipennis*. If Richards' description of *Isodontia bastiniana* is accurate his species is distinct from *nigrocoerulea*. Richards states that the holotype of *bastiniana* lacks the prominent transverse bands of pale cilia found on the gastral sternites of *nigrocoerulea*. In addition, the flagellomeres of *bastiniana* do not possess the spicules found on the flagellomeres of *nigrocoerulea*. A possibility

exists then that *cyanipennis* and *bastiniana* are synonymous and distinct from *nigrocoerulea*. All of the blue *Isodontia* that I have seen from the type locality of *cyanipennis*, Cayenne, are *nigrocoerulea* however, and it seems best to follow van der Vecht's interpretation of *cyanipennis* for the present.

In his original description, Taschenberg indicated that he was describing a male and a female, and the two syntypes sent to me are so labeled but both are females.

The type of *I. bipunctata* Rohwer is identical with Taschenberg's *nigrocoerulea*.

#### SPHEX TEPANECUS Saussure

*Sphex tepanecus* Saussure, 1867. Reise der Ost. Freg. Novara, Zool., 2:41. (holotype ♂, "Mextill," Museum d'histoire Naturelle, Geneva).

*Sphex mexicana* Taschenberg, 1869. Zeitsch. Ges. Naturw. Halle, 34:416 (holotype ♂, Mexico).

Examination of Taschenberg's type proves that Kohl (1890) was correct in synonymizing *mexicanus* with *tepanecus* Saussure. Specimens of *tepanecus* that I have studied from Arizona and Texas differ from the type of *mexicanus* only slightly. The legs are completely black on the type but the United States examples have the front femora reddish brown beneath.

#### SPHEX ARGENTINUS Taschenberg

*Sphex argentina* Taschenberg, 1869. Zeitsch. Ges. Naturw. Halle, 34:417 (lectotype ♂, Mendoza, Argentina; paralectotype ♀, Rozario, Argentina, present designation).

Willink (1951) correctly interpreted this species.

#### SPHEX MELANOPUS Dahlbom

*Sphex melanopa* Dahlbom, 1843, Hymenoptera Europaea, 1:27 (holotype ♂, Brazil, Zoologisches Museum, Humboldt Universitat, Berlin.)

*Sphex proxima* Smith, 1856. Cat. Hym. Ins. Coll. Brit. Mus., 4:258 (holotype ♀, Brazil, British Museum Natural History).

*Sphex ruficauda* Taschenberg 1869. Zeitsch. Ges. Naturw. Halle, 34:418 (holotype ♂, "Amer. Merid.")

Both Kohl (1895) and Fernald (1931) studied Dahlbom's type of *melanopus* and arrived at the above synonymy. Dahlbom mentioned that his type was in the collection at Lund, but Fernald could not locate it there, and stated, as did Kohl, that it was in Berlin. Fernald also studied Smith's type of *proxima*.

This species exhibits a north-south clinal variation in the amount and color of thoracic pubescence. On specimens from Brazil, the appressed pubescence is copper or tarnished silver and is confined to a small spot behind the pronotal lobe and

above the meso- and meta-coxa, and to the scutal furrows and also a small spot at the side of the petiole socket. In these specimens the erect pubescence is dirty white. The area covered by the appressed pubescence increases on specimens found further north. In Venezuelan examples the pronotal lobe is covered with appressed silver pubescence and the posterior portion of the propodeum is entirely covered by appressed silver pubescence. Erect thoracic hair on these specimens is golden. In Panamanian examples the propodeum is completely covered and the thorax is extensively covered by appressed golden pubescence. Taschenberg's type of *ruficauda* agrees most closely with males from Venezuela.

#### SPHEX DORSALIS Lepeletier

*Sphex dorsalis* Lepeletier, 1845. Hist. Nat. Insect. Hym., 3:347 (holotype ♂, Cayenne, French Guiana, Museo di Zoologia, Universita di Torino (Turin)).

*Sphex singularis* Smith, 1856. Cat. Hym. Insect. Brit. Mus., 4:261. (holotype ♂, Honduras, British Museum Natural History).

*Sphex micans* Taschenberg, 1869. Zeitsch. Ges. Naturw. Halle, 34:419 (lectotype ♀ and paralectotype ♀, Parana, Argentina, present designation).

Of the five female specimens sent to me for study, only two are clearly Taschenberg syntypes of *micans*. Taschenberg listed Parana, Mendoza, and Rio de Janeiro as type localities for the five females he described. Two of the specimens before me lack labels and a third is labeled Rozario. I am restricting the designation of lectotype to one of the two females labeled Parana.

#### LITERATURE CITED

FERNALD, H. T.

1931. Notes on some American Sphecinae. Ann. Ent. Soc. Amer., 24(3):439-450.

KOHL, F.F.

1890. Die Hymenopterengruppe der Sphecinen, I. Monographie der Naturlichen gattung Sphex. Ann. k.k. Naturhist. Hofmus. Wien, 5:77-194, 317-461.

1895. Zur monographie der naturlichen gattung Sphex Linne. Ann. k.k. Naturhist. Hofmus Wien, 10:42-74.

RICHARDS, O. W.

1937. Results of the Oxford University Expedition to British Guiana, 1929. Hymenoptera, Sphecidae and Bembicidae. Trans. Royal Ent. Soc. London, 86(7):101-118.

VAN DER VECHT, J.

1961. Hymenoptera Sphecoidea Fabriciana. Zool. Verh., no. 48:1-85.

WILLINK A.

1951. Las especies Argentinas y Chilenas de Chlorionini. Acta Zool. Lilloana, 11:53-225.